

A comment on a paper by Carot et al.

Alan Barnes

Computer Science Group, School of Engineering and Applied Science,
Aston University, Aston Triangle, Birmingham B4 7ET, UK [†]

PACS number(s): 04.20

Submitted to Class. Quantum Grav.

Date: 7 February 2008

Abstract. In a recent paper Carot et al. considered carefully the definition of cylindrical symmetry as a specialisation of the case of axial symmetry. One of their propositions states that if there is a second Killing vector, which together with the one generating the axial symmetry, forms the basis of a two-dimensional Lie algebra, then the two Killing vectors must commute, thus generating an Abelian group. In this comment a similar result, valid under considerably weaker assumptions, is recalled: any two-dimensional Lie transformation group which contains a one-dimensional subgroup whose orbits are circles, must be Abelian. The method used to prove this result is extended to apply to three-dimensional Lie transformation groups. It is shown that the existence of a one-dimensional subgroup with closed orbits restricts the Bianchi type of the associated Lie algebra to be I (Abelian), II, III, VII_{q=0}, VIII or IX. The relationship between the present approach and that of the original paper is discussed.

[†] E-Mail: barnesa@aston.ac.uk Fax: +44 121 333 6215

1. Introduction

We begin by recalling a few relevant definitions and theorems. Following Carter (1970) a spacetime $\mathcal{M}$ is said to have cyclical symmetry if and only if the metric is invariant under the effective smooth action $SO(2) \times \mathcal{M} \rightarrow \mathcal{M}$ of the one-parameter cyclic group $SO(2)$. If, in a cyclically symmetric spacetime, the set of fixed points of the isometry is non-empty then the set of fixed points is an autoparallel time-like two-surface (Mars and Senovilla, 1993) and the spacetime is said to be axially symmetric. Mars and Senovilla also show that the Killing vector is spacelike in the neighbourhood of the axis and satisfies the elementary flatness condition on the axis.

In a recent paper Carot, Senovilla and Vera (1999) consider in detail the definition of cylindrical symmetry; this paper will be referred to below as CSV. Proposition 2 of CSV states that in an axially symmetric spacetime, if there is another Killing vector which, with the axial Killing vector generates a two-dimensional isometry group, then the two Killing vectors commute. Thus the isometry group is Abelian. A similar result for stationary axisymmetric spacetimes was proved by Carter (1970). Both proofs rely heavily on the existence of an axis where the axial Killing vector vanishes. CSV then use their result as the basis of a straightforward definition of a cylindrical symmetry without the additional assumptions that are often made namely: that the group is Abelian and that an elementary flatness condition holds at the axis. Thus a spacetime is cylindrically symmetric if it admits a two-dimensional isometry group with spacelike orbits (i.e. a G_2 acting on S_2) containing an axial symmetry.

Although the assumption of the existence of an axis is reasonable in many circumstances, there are physically realistic situations where an axis may not exist: the axis may be singular and so not part of the manifold, or the topology of the manifold may be such that no axis exists. An example of the latter situation is a torus embedded in a 3-dimensional flat space; the torus is cyclically symmetric but the Killing vector field generating the rotation does not vanish anywhere on the surface. It is interesting therefore to consider whether a result of the same ilk as Proposition 2 of CSV is valid for cyclically symmetric spacetimes.

2. Cyclically symmetric manifolds admitting a G_2

Suppose $\mathcal{M}$ is cyclically symmetric and suppose X_0 is the Killing vector field generating the isometry. We restrict attention to the open submanifold $\mathcal{N}$ of $\mathcal{M}$ on which X_0 is non-zero. The orbit each point of $\mathcal{N}$ under the cyclic symmetry is a circle and these circles are the integral curves of the Killing vector field X_0 . Let ϕ be a circular coordinate running from 0 to 2π which labels the elements of $SO(2)$ in the usual way. Then we can introduce a system of coordinates x^i with $i = 1 \dots n$ and $x^1 = \phi$ adapted to X_0 such that $X_0 = \partial_\phi$. These coordinates are determined only up to transformations of the form

$$\tilde{\phi} = \phi + f(x^\nu) \quad \tilde{x}^\mu = g^\mu(x^\nu)$$

where f and g^μ are smooth functions and where Greek indices take values in the range $2 \dots n$.

Suppose now that the isometry group of $\mathcal{M}$ contains a two-dimensional subgroup G_2 and that the Killing vector field X_1 together with X_0 forms a basis of the associated Lie algebra with commutation relations $[X_0, X_1] = aX_0 + bX_1$, say, where a and b are constants. In an adapted coordinate system the commutation relations reduce to

$$\frac{\partial X_1^\mu}{\partial \phi} = bX_1^\mu \quad \frac{\partial X_1^1}{\partial \phi} = a + bX_1^1$$

A straightforward integration gives

$$\begin{aligned} X_1^\mu &= B^\mu(x^\nu) e^{b\phi} & X_1^1 &= A(x^\nu) e^{b\phi} - a/b & \text{for } b \neq 0 \\ X_1^\mu &= B^\mu(x^\nu) & X_1^1 &= A(x^\nu) + a\phi & \text{for } b = 0 \end{aligned}$$

where A and B^μ are arbitrary functions of integration. The solutions of these equations must be periodic in ϕ with period 2π if X_1 is to be single-valued on $\mathcal{N}$. Thus both a and b must vanish and the subgroup G_2 must be Abelian.

Note that the dimensionality of the manifold, the existence of a metric and the fact that transformation group is an isometry group are not used in the proof; thus it has been proved that any two-dimensional Lie transformation group which acts on an n -dimensional manifold $\mathcal{M}$ and which contains a one-dimensional subgroup acting cyclically on $\mathcal{M}$ must be Abelian. This remarkably simple and general result is not new[‡], but is perhaps not widely known.

3. Cyclically symmetric manifolds admitting a G_3

Suppose now that the manifold $\mathcal{M}$ admits a three-dimensional Lie group G_3 of transformations which contains a one-dimensional subgroup, generated by the vector field X_0 , acting cyclically on $\mathcal{M}$. Let X_1 and X_2 be vector fields on $\mathcal{M}$ which together with X_0 form a basis of the associated Lie algebra. Two cases arise: either the Lie algebra admits a two-dimensional subalgebra containing X_0 or there is no such subalgebra.

In the former case, by the result of the previous section the subalgebra is Abelian. We assume without loss of generality that X_0 and X_1 form a basis of this subalgebra. Hence the commutation relations can be written in the form

$$[X_0, X_1] = 0 \quad [X_0, X_2] = aX_0 + bX_1 + cX_2$$

where a , b and c are constants. In terms of a coordinate system adapted to X_0 these commutators reduce to the differential equations

$$\frac{\partial X_1^i}{\partial \phi} = 0 \quad \frac{\partial X_2^i}{\partial \phi} = a\delta_1^i + bX_1^i + cX_2^i$$

[‡] The author remembers seeing this result presented as a footnote in a paper around 30 years ago, but is unable to locate the precise reference.

A straightforward argument similar to that in the previous section shows that if the solutions are to be periodic in ϕ , we must have $a = b = c = 0$. Thus X_0 commutes with the other two basis vectors X_1 and X_2 . The remaining commutator takes the form

$$[X_1 X_2] = dX_0 + eX_1 + fX_2$$

where d , e and f are constants. Three algebraically distinct cases arise. If $d = e = f = 0$ the group is Abelian (Bianchi type I). If $e = f = 0$ but $d \neq 0$, after a rescaling of X_1 and/or X_2 , we may set $d = 1$. The commutation relations now are

$$[X_0 X_1] = 0 \quad [X_0 X_2] = 0 \quad [X_1 X_2] = X_0$$

which is the canonical form for an algebra of Bianchi type II (see for example Petrov, 1969). If e and f are not both zero, by a change of basis which leaves X_0 fixed, we can set $d = e = 0$ and $f = 1$. Thus the commutators are

$$[X_0 X_1] = 0 \quad [X_0 X_2] = 0 \quad [X_1 X_2] = X_2$$

which is (essentially) the canonical form for an algebra of Bianchi type III.

If the Lie algebra has no two-dimensional subalgebra containing X_0 then, without loss of generality, we may choose the basis vectors X_1 and X_2 such that the commutation relations become

$$[X_0 X_1] = X_2 \quad [X_0 X_2] = aX_0 + bX_1 + cX_2$$

where a , b and c are constants. In terms of a coordinate system adapted to X_0 these reduce to

$$\frac{\partial X_1^i}{\partial \phi} = X_2^i \quad \frac{\partial X_2^i}{\partial \phi} = a\delta_1^i + bX_1^i + cX_2^i$$

From these equations it is easy to deduce that

$$\frac{\partial^2 X_2^i}{\partial \phi^2} = bX_2^i + c\frac{\partial X_2^i}{\partial \phi}$$

Thus for solutions periodic in ϕ with period 2π , we must have $c = 0$ and $b = -n^2$ for some positive integer n . By a redefinition of the basis vector $\tilde{X}_1 = X_1 + a/bX_0$, we can also set $a = 0$ so that the commutation relations become

$$[X_0 X_1] = X_2 \quad [X_0 X_2] = -n^2 X_1 \quad [X_1 X_2] = dX_0 + eX_1 + fX_2$$

where d , e and f are constants. In a coordinate system adapted to X_0 the basis vectors are given by

$$\begin{aligned} X_0^i &= \delta_\phi^i \\ X_1^i &= 1/n(A^i(x^\nu) \sin n\phi - B^i(x^\nu) \cos n\phi) \\ X_2^i &= A^i(x^\nu) \cos n\phi + B^i(x^\nu) \sin n\phi \end{aligned}$$

where A^i and B^i are functions of integration.

The commutators must, of course, satisfy the Jacobi identity

$$[X_0 [X_1 X_2]] + [X_1 [X_2 X_0]] + [X_2 [X_0 X_1]] = 0$$

which implies that

$$[X_0 (dX_0 + eX_1 + fX_2)] = eX_2 - n^2 fX_1 = 0$$

Thus $e = f = 0$. Three algebraically distinct cases arise. If $d = 0$ the group is of Bianchi type VII _{$q=0$} which is, of course, the isometry group of the Euclidean plane. Note that the standard form of the commutation relations for Bianchi type VII _{$q=0$} is obtained by means of the change of basis

$$\tilde{X}_0 = 1/nX_0 \quad \tilde{X}_1 = nX_1 \quad \tilde{X}_2 = X_2$$

For $d < 0$ and $d > 0$ the Bianchi type is VIII and IX respectively; which are of course the symmetry groups of Lobachevskii plane and two-sphere respectively. The change of basis required to reduce the commutation relations to the canonical form for Bianchi types VIII and IX is

$$\tilde{X}_0 = 1/nX_0 \quad \tilde{X}_1 = n/\epsilon X_1 \quad \tilde{X}_2 = 1/\epsilon X_2$$

where $\epsilon = n\sqrt{|d|}$.

The methods above can be applied to the case of four (and higher) dimensional Lie groups containing a one-dimensional cyclic subgroup. Again the allowed forms of the commutation relations are considerably restricted. These results will be presented elsewhere.

4. A correction

Senovilla (1999) has pointed out an error in the statement of Theorem 3 of CSV. At his request a corrected statement of the theorem is presented here.

Given a G_3 on T_3 group that contains an orthogonally transitive Abelian G_2 subgroup generated by an axial $\vec{\xi}$ (such that its set of fixed points is non-empty) and $\vec{\eta}$, and a third integrable timelike Killing vector field $\vec{\zeta}$, then it follows that: if G_3 is the maximal isometry group, then it must be Abelian; if G_3 is non-Abelian, then it is (locally) contained in a G_4 on T_3 , and $\vec{\xi}$ and $\vec{\zeta}$ generate an orthogonally transitive subgroup G_2 on T_2 .

The proof follows easily from the remarks immediately preceding the statement of the theorem in the original paper.

5. Summary

Low-dimensional Lie transformation groups which act on an n -dimensional manifold $\mathcal{M}$ and which contain a one-dimensional subgroup acting cyclically on $\mathcal{M}$ have been considered. For the two-dimensional case the group must be Abelian and for the three-dimensional case the Bianchi type of the group is restricted: types IV, V and VI cannot occur and only the subclass $\text{VII}_{q=0}$ of type VII is allowed. As far as the author is aware the results for the three-dimensional case are new.

At first sight the restrictions in the three-dimensional case may not seem too severe: only three of the nine Bianchi types are excluded whilst a fourth type is restricted. However, it should be pointed out that Bianchi types VI and VII both involve an arbitrary parameter and so each contain an infinite number of algebraically distinct types. Those in type VI are excluded completely and for type VII only a single case survives. Moreover for most of the types which *are* permitted the structure of the Lie algebra is ‘aligned’ in some way relative to X_0 . For example, for Bianchi type II, X_0 is also the generator of the first derived subalgebra and for Bianchi types II and III, X_0 lies in the centre of the algebra.

At first sight too, the result presented in section 2 seems considerably stronger than Proposition 2 of CSV; however this is only partially true. Certainly the results presented do not require the existence of an axis and they apply to any Lie transformation group. So *a fortiori* the results apply to isometries, conformal motions, affine and projective collineations etc. whereas CSV’s results apply only to isometries (although some of the results, at least, may be extended to conformal motions). However the results presented here require that the transformations are defined everywhere on the manifold and so they say nothing about the structure of groups of *local* transformations. CSV’s results apply to local isometries in the neighbourhood of the axis and so in this sense are stronger than the results presented above. Of course, if we assume that the isometries are globally well-defined and that an axis of fixed points exists, it may well be possible to derive stronger restrictions on the allowed group structure using a synthesis of the two methods. This is currently under investigation.

References

- Carot J, Senovilla J M M and Vera R *Class. Quantum Grav.* **16** 3025–34
 Carter B 1970 *Commun. math. Phys.* **17** 233–8
 Mars M and Senovilla J M M 1993 *Class. Quantum Grav.* **10** 1633–47
 Petrov A Z 1969 *Einstein Spaces* Pergamon Press, Oxford, p. 63
 Senovilla J M M 1999 *Private Communication*